

Analysis of an irregular boundary layer behavior for the steady state flow of a Boussinesq fluid

Christos Sourdis

University of Turin

'This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 609402 - 2020 researchers: Train to Move (T2M)'



Turcotte, Spence and Bau *Int. J. Heat Mass Transfer* (1982) considered the vertical flow of an internally heated Boussinesq fluid in a vertical channel with viscous dissipation and pressure work.

Boundary value problem for the steady state flow:

$$\begin{cases} 2u'' = u^2 - A(1 - x^2), & x \in (-1, 1), \\ u(-1) = u(1) = 0, \end{cases}$$

A represents the (non-dimensional) heat addition

u is the velocity

x is the scaled position

$[-1, 1]$ is the horizontal cross section of the vertical channel

$$\begin{cases} 2u'' = u^2 - A(1 - x^2), & x \in (-1, 1), \\ u(-1) = u(1) = 0, \end{cases}$$

Numerical simulations indicated that the number of (even) solutions diverges as $A \rightarrow \infty$.

Setting

$$\varepsilon^2 = \frac{2}{\sqrt{A}} \rightarrow 0 \text{ and } v = \frac{1}{\sqrt{A}}u,$$


the problem is equivalent to the singular perturbation problem:

$$\begin{cases} \varepsilon^2 v'' = v^2 - (1 - x^2), & x \in (-1, 1), \\ v(-1) = v(1) = 0. \end{cases}$$


The $\varepsilon = 0$ limit problem (outer problem)

$$\begin{cases} 0 = v^2 - (1 - x^2), & x \in (-1, 1), \\ v(-1) = v(1) = 0. \end{cases}$$

Two continuous solutions : $v_0 = \pm\sqrt{1 - x^2}$.



It is natural to expect that there exist solutions $v_\varepsilon \rightarrow v_0$ uniformly on $[-1,1]$ as $\varepsilon \rightarrow 0$.



However, a more complete formal analysis is required, with special attention at the boundary where such solutions should develop **irregular boundary layer behavior**.

Blow-up problem

Near the boundary $x = -1$, say for $x \in [-1, -1 + \delta]$, a standard matching asymptotic analysis **Turcotte, Spence and Bau** *Int. J. Heat Mass Transfer* (1982) predicts that


$$v_\varepsilon(x) \sim \varepsilon^{2/5} 2^{2/5} Y(s), \quad s = 2^{1/5}(x + 1)/\varepsilon^{4/5}$$

$$\begin{cases} y'' = y^2 - s, & s > 0, \\ y(0) = 0, & y - \sqrt{s} \rightarrow 0 \text{ as } s \rightarrow \infty. \end{cases}$$


The above differential equation is integrable, known as the **Painlevé-I transcendent**.

Holmes-Spence (1984) used dynamical systems techniques to find a unique increasing solution Y_+ and at least one solution Y_- with exactly one local minimum (see also **Dancer-Yan (2005)** for variational methods).

Hastings-Troy (1989) showed that these two are the only solutions (their proof relied on some four decimal point numerical calculations!)



$$\begin{cases} y'' = y^2 - s, & s > 0, \\ y(0) = 0, & y - \sqrt{s} \rightarrow 0 \text{ as } s \rightarrow \infty. \end{cases}$$



Nondegeneracy

$$-\phi'' + 2Y_{\pm}(s)\phi = 0, \quad s > 0, \quad \phi(0) = 0, \quad \phi \in L^{\infty}(0, \infty) \Rightarrow \phi \equiv 0.$$

Proposition [Sourdis 2015]

$$-\phi'' + 2Y_-(s)\phi = 0, \quad s > 0, \quad \phi(0) = 0, \quad \phi \in L^\infty(0, \infty) \Rightarrow \phi \equiv 0.$$

Idea of the proof

$$y'' = y^2 - s; \quad y(0) = 0, \quad y(s) \sim \sqrt{s}, \quad s \rightarrow \infty.$$

We know that there is a unique positive solution Ψ_+ , with $\Psi'_+ > 0$. Following **Dancer-Yan (2005)**, we search a sign changing solution $Y_- < Y_+$ as $Y_- = Y_+ - w$, where w is a positive solution of the '**Nonlinear Schrödinger Equation**'

$$w'' - V(s)w + w^2 = 0, \quad s > 0, \quad w(0) = 0, \quad w(\infty) = 0,$$

with potential $V(s) = 2Y_+(s)$

$$V(0) = 0, \quad V' > 0, \quad V \sim 2\sqrt{s} \text{ as } s \rightarrow \infty$$

w is a mountain pass solution to the above problem

$$w'' - V(s)w + w^2 = 0, \quad s > 0, \quad w(0) = 0, \quad w(\infty) = 0,$$

$$V(0) = 0, \quad V' > 0, \quad V \sim 2\sqrt{s} \text{ as } s \rightarrow \infty$$

Felmer-Martinez-Tanaka (2008) Uniqueness and non-degeneracy result includes:

$$w'' + (\nu/s)w' - V(s)w + w^2 = 0, \quad s > a, \quad w(a) = 0 \quad (a > 0), \quad w(\infty) = 0,$$

$$\nu > 0, \quad 0 < \inf V < \sup V < \infty, \quad V' \geq 0$$

Modelled after $\Delta u - u + u^p = 0, \quad x \in \mathbb{R}^n, \quad |x| > a \quad (\nu = n - 1)$.

Long history: Coffman, Kwong, McLeod-Serrin, Peletier-Serrin, Ni-Takagi, Tang ...

Outer solution: $\sqrt{1 - x^2}$

Near $x = -1$, two possible inner solutions:

$$\varepsilon^{2/5} 2^{2/5} Y_{\pm} \left(2^{1/5} (x + 1) / \varepsilon^{4/5} \right)$$

Glue them together and then show that the resulting global approximate solution can be perturbed to a genuine solution

PDE approach: Sourdís-Fife (2007), Karali-Sourdís (2012)

Blow-up approach to geometric singular perturbation theory: Krupa-Szmolyan (2001), Schechter (2004)
Schechter-Sourdís (2009)

Theorem [Sourdis (2015)]

$$\begin{cases} \varepsilon^2 v'' = v^2 - (1 - x^2), & x \in I = (-1, 1), \\ v(-1) = v(1) = 0. \end{cases}$$

For small $\varepsilon > 0$ has four solutions such that


$$\|v_\varepsilon - \sqrt{1 - x^2}\|_{L^\infty(-1,1)} \leq C\varepsilon^{2/5}$$

$$v_\varepsilon(x) = 2^{2/5} \varepsilon^{2/5} Y_i \left(2^{1/5} \text{dist}(x, \partial I) / \varepsilon^{4/5} \right) + O(\varepsilon^{6/5}),$$

for $\text{dist}(x, \partial I) \leq C\varepsilon^{4/5}$, $i = \pm$.

Moreover, they are **nondegenerate**:

$$-\varepsilon^2 \phi'' + 2v_\varepsilon \phi = f, \quad \phi(-1) = \phi(1) = 0 \Rightarrow \|\phi\|_{L^\infty} \leq C\varepsilon^{-2/5} \|f\|_{L^\infty}$$



Morse index 1 solution (see also **Dancer-Yan** (2005))


Dancer-Yan (2005) gave the first proof of the **Lazer-McKenna conjecture** from 1983 in the PDE case by establishing that the number of solutions diverges as $\varepsilon \rightarrow 0$ for the singular perturbation problem:

$$\varepsilon^2 \Delta u = |u|^p - \phi_1(x), \quad x \in \Omega \subset \mathbb{R}^n; \quad u = 0, \quad x \in \partial\Omega$$


$\phi_1 > 0$ eigenfunction associated to $\lambda_1(\Omega)$ ($\partial_\nu \phi_1 < 0$ on $\partial\Omega$)

$1 < p < (n + 2)/(n - 2)$ if $n \geq 3$, $p > 1$ if $n = 1, 2$.

Outer solution: $\phi_1^{1/p}$



$$\varepsilon^2 \Delta w - 2u_\varepsilon(x)w + w^2 = 0, \quad x \in \Omega, \quad w > 0, \quad w = 0 \quad x \in \partial\Omega$$



Small peaks along manifolds: [Byeon-Oshita \(2015\)](#)

THANK YOU FOR YOUR ATTENTION